arXiv:1411.6084v3 [math.AG] 10 Dec 2014

ON THE CUT-AND-PASTE PROPERTY OF ALGEBRAIC VARIETIES

ILYA KARZHEMANOV

Abstract. We show that the above-named property (after M. Larsen and V. Lunts) does not hold in general.

1. Introduction

1.1. Fix some ground field $\mathbf{k} \subseteq \mathbb{C}$. Consider the ring $K_0(\mathrm{Var}_{\mathbf{k}})$ of $\mathbf{k}$-varieties, generated (over $\mathbb{Z}$) by the isomorphism classes $[X]$ of various quasi-projective such X, with multiplication given by $[X] \cdot [Y] := [X \times_{\mathbf{k}} Y]$ (for all Y) and addition $[X] + [Y]$ being just the formal sum subject to the relation $[X] = [X \setminus Y] + [Y]$ whenever $Y \subseteq X$ is a closed subset. In particular, the class of a point can be identified with the unity $1 \in K_0(\mathrm{Var}_{\mathbf{k}})$, so that $[\mathbb{A}^1_{\mathbf{k}} \setminus \{0\}] = \mathbb{L} - 1$ for $[\mathbb{A}^1_{\mathbf{k}}] := \mathbb{L}$. We will suppress the reference to $\mathbf{k}$ in what follows for notations' simplicity.

The ring $K_0(\mathrm{Var})$, or more precisely, *motivic measures* with values in $K_0(\mathrm{Var})$, was used in [8] for example to show (via the celebrated *motivic integration*) that birationally isomorphic Calabi-Yau manifolds have equal Hodge numbers (see [3], [4], [11], [12], [9], [10] and [5] for other results, applications and references). Despite this success, however, the structure of $K_0(\mathrm{Var})$ is still poorly understood, although it is known that $K_0(\mathrm{Var})$ is not a domain (see [13]) and one has an explicit description of "conical" subrings in $K_0(\mathrm{Var})$ (see [7]); see also [9] and [1] for a description of quotients of $K_0(\mathrm{Var})$ by some ideals related to $\mathbb{L}$ (note that it is still not known whether $\mathbb{L}$ is a 0-divisor in $K_0(\mathrm{Var})$). The present paper provides another (tiny) contribution to this interesting subject.

Namely, given two varieties X and Y let's say (following [9]) that they satisfy the *cut-and-paste property* if $[X] = [Y]$ in $K_0(\mathrm{Var})$ and additionally $X = \coprod_{i=1}^{k} X_i$ (resp. $Y = \coprod_{i=1}^{k} Y_i$) for some $k \geq 1$ and constructible subsets $X_i \subseteq X$ (resp. $Y_i \subseteq Y$), with $X_i \simeq Y_i$ for all i.

Theorem 1.2. *There exist two smooth projective varieties X and Y which violate the cut-and-paste property.*[1)]

In order to prove Theorem 1.2 it suffices to exhibit such X and Y, not birational to each other, yet satisfying $[X] = [Y]$. Specifically, the idea is to take *birationally rigid* X and Y (provided, say, by the paper [14]), and then show that $[X] = [Y]$. For the latter, we want both X, Y to be "simple", e.g. fibred over $\mathbb{P}^1$ onto surfaces all having the same class in $K_0(\mathrm{Var})$ (this is also inline with [14]). The equality $[X] = [Y]$, for certain X and Y, can then be established via direct technical (though elementary) argument (see **2.1**, **2.5** below).

[1)] This also answers Question (b) in [6, **3.G'''**] at negative due to the results from [2].

2. Proof of Theorem 1.2

2.1. Fix two cubic forms $G := G(x_0, \dots, x_3)$ and $F := F(x_0, \dots, x_3)$ over $\mathbf{k}$. We will assume G generic and the equation $F = 0$ define a cubic surface (in $\mathbb{P}^3$) with an ordinary double point as the only singularity. We also choose F generic among such forms.

Consider the locus $X \subset \mathbb{P}^3 \times \mathbb{P}^1$ given by an equation

$$\alpha(t_0, t_1)G + \beta(t_0, t_1)F = 0 \tag{2.2}$$

of bidegree $(3, m)$ for G, F as above and some $m \geq 3$, with x_i (resp. t_i) being projective coordinates on $\mathbb{P}^3$ (resp. on $\mathbb{P}^1$). We also assume both forms α, β to be generic.

Lemma 2.3. *X is a smooth 3-fold.*

Proof. Indeed, by Bertini theorem applied to the linear system of divisors (2.2), all possible singularities of X can lie only on the surface $\mathfrak{B} := (G = F = 0) \subset \mathbb{P}^3 \times \mathbb{P}^1$. This surface is smooth as being isomorphic to $\mathbb{P}^1\times$ the smooth curve $(G = F = 0) \subset \mathbb{P}^3$. In particular, locally analytically near every point on $\mathfrak{B}$ we may assume both G, F to be some local coordinates. The claim now easily follows by taking partial derivatives of (2.2) w. r. t. t_i and G, F. □

Further, X carries a natural fibration $X \longrightarrow \mathbb{P}^1$ on cubic surfaces induced by the projection of $\mathbb{P}^3 \times \mathbb{P}^1$ onto the second factor. We will refer to X as a *pencil* (of cubic surfaces).

Let $y_0 := t_0^m, y_1 := t_0^{m-1}t_1, \dots, y_m := t_1^m$ be the monomials entering α and β with some $\mathbf{k}$-coefficients $\alpha_0, \dots, \alpha_m$ and $\beta_0, \dots, \beta_m$, respectively. We may regard y_i as new transcendental variables. Consider the locus $\mathcal{X} \subset \mathbb{P}^3 \times \mathbb{P}^m \times \mathbb{A}^1$ given by equation

$$L_\alpha G + (L_\beta + \lambda)F = 0,$$

where $\mathbb{A}^1 = \operatorname{Spec} \mathbf{k}[\lambda]$ and the linear forms $L_\alpha := L_\alpha(y_0, \dots, y_m), L_\beta := L_\beta(y_0, \dots, y_m)$ are obtained from α, β, respectively, via replacing each monomial with the corresponding y_i, i.e., $L_\alpha = \sum \alpha_i y_i$ and $L_\beta = \sum \beta_i y_i$.

Put $t_0 := 1$ (resp. $y_0 := 1$) and consider the open loci

$$X_0 := X \cap (t_0 = 1) \setminus (G = F = 0) \subset X$$

and

$$\mathcal{X}_0 := \mathcal{X} \cap (y_0 = 1) \setminus (G = F = 0) \subset \mathcal{X}.$$

Notice that $X_0 \subset \mathcal{X}_0 \times \mathbb{A}^1$ as a closed subset, with (tautological) equations $y_1 = t, \dots, y_m = t^m$, where $t := t_1$. More precisely, we have

Lemma 2.4. *$X_0 \times \mathbb{A}^{m+1} \simeq \mathcal{X}_0 \times \mathbb{A}^1$ and $(X_0 \times \mathbb{A}^{m+1}) \cap (y_i = 0) \simeq (\mathcal{X}_0 \times \mathbb{A}^1) \cap (y_i = t)$ for all i, where $\mathbb{A}^{m+1}$ (resp. $\mathbb{A}^1$) is provided with affine coordinates $y_1, \dots, y_m, \lambda$ (resp. t).*

Proof. Consider $\mathbb{A}^2 = \operatorname{Spec} \mathbf{k}[\lambda, z]$ and identify $\mathcal{X}_0 \times \mathbb{A}^1$ with the closed subset $(y_1 = z) \subset \mathcal{X}_0 \times \mathbb{A}^2$. Let's construct an isomorphism $\varphi : X_0 \times \mathbb{A}^{m+1} \xrightarrow{\sim} \mathcal{X}_0 \times \mathbb{A}^1$ as follows:

$$[x_0 : \ldots : x_3] \times t \times (y_1, \ldots, y_m, \lambda) \mapsto [x_0 : \ldots : x_3] \times (y'_1, y'_2, \ldots, y'_m, \lambda') \times t \times y_1,$$
$$y'_1 := y_1 + t + \lambda,\ y'_2 := y_2 + t^2,\ \ldots,\ y'_m := y_m + t^m,\ \lambda' := (-\beta_1 - \alpha_1 \tfrac{G(x)}{F(x)})\lambda - L_\beta(y) + \beta_0 - \tfrac{G(x)}{F(x)}(L_\alpha(y) - \alpha_0),$$

where $x := (x_0, \ldots, x_3)$ and similarly for y.[2)]

The obtained morphism $\varphi : X_0 \times \mathbb{A}^{m+1} \longrightarrow \mathcal{X}_0 \times \mathbb{A}^1$ is easily seen to be dominant (by dimension count) and one-to-one (for it is this on the fibers = cubic surfaces of the form (2.2)). Indeed, we may fix x_i (all i), y_1 and t. Then φ reduces to an *affine* map on $x \times t \times \mathbb{A}^m$ sending $(y_2, \ldots, y_m, \lambda)$ to $(y'_2, \ldots, y'_m, \lambda')$ as is indicated above. (The corresponding equation $L_\alpha(y')G(x) + (L_\beta(y') + \lambda')F(x) = 0$ turns into $y'_1 = y_1 + t + \lambda$.) It's now immediate from the construction that $\varphi\big|_{x \times t \times \mathbb{A}^m}$ is an isomorphism.

Thus φ is an isomorphism. The last assertion of lemma is evident. □

2.5. Consider another locus $\tilde{X} \subset \mathbb{P}^3 \times \mathbb{P}^1$, defined similarly as X with the same G, F, but having other (still generic) degree m forms $\tilde{\alpha}(t_0, t_1), \tilde{\beta}(t_0, t_1)$. All the previous gadgets such as $\mathcal{X}, X_0$, etc. are defined verbatim for $\tilde{X}$, and we will distinguish them (from those for X) by simply putting extra $\tilde{}$.

One may assume w. l. o. g. the fiber over the point $[0 : 1]$ for $X \longrightarrow \mathbb{P}^1$ (resp. for $\tilde{X} \longrightarrow \mathbb{P}^1$) to be *smooth*. Further, observe that both X and $\tilde{X}$ have the same number of singular fibers, all being of that type as the surface $(F = 0)$ (cf. the beginning of **2.1**). Indeed, the pencils $X, \tilde{X}$ correspond to smooth rational curves $(C, \tilde{C}$, say) in the space of all cubics in $\mathbb{P}^3$, with F corresponding to generic point in the (closed) locus Σ of all singular such cubics. But C and $\tilde{C}$ can be chosen to intersect Σ only at generic points (for instance, use the parameter count and generality of $X, \tilde{X}$), and the claim follows as $\chi_{\text{top}}(X) = \chi_{\text{top}}(\tilde{X}) = 18 - s$ for $s =$ the number of singular fibers in each of $X, \tilde{X}$.[3)]

Remark 2.6. More precisely, letting $m := 1$ in the definition of X (see (2.2)), one obtains the blowup $\operatorname{Bl}_Z(\mathbb{P}^3)$ of $\mathbb{P}^3$ at the curve $Z := (G = F = 0)$. This immediately gives $\chi_{\text{top}} = -14$ for such X. In general, there is a finite morphism $X \longrightarrow \operatorname{Bl}_Z(\mathbb{P}^3)$ of degree m and with only *simple* branch locus, concentrated on smooth fibers. Then we get $\chi_{\text{top}}(X) = m(-14 - 9(2m-2)) + 9(2m-2)(m-1) = -14m - 9(2m-2) = -32m + 18$ and thus $s = 32m$.

With this setup, we argue as follows:

Proposition 2.7. *$[X] \cdot \mathbb{L}^k = [\tilde{X}] \cdot \mathbb{L}^k$ for all $1 \le k \le m+1$. Furthermore, there is an isomorphism $X_0 \times \mathbb{A}^{m+1} \simeq \tilde{X}_0 \times \mathbb{A}^{m+1}$, and similarly for m instead of $m+1$.*

Proof. Firstly, since

$$[X] = [X_0] + [\text{the fiber over } [0:1]] + [Z] \cdot \mathbb{L}$$

[2)] Heuristically, we "spread out" the locus $X_0 \subset \mathcal{X}_0 \times \mathbb{A}^1$ over the whole $\mathcal{X}_0 \times \mathbb{A}^1$, shifting it by y_i and λ.

[3)] In the latter argument, we have used that χ_{top} (a. k. a. $\chi_{\text{ét}}$) of the cubic surface $(G = 0)$ equals 9, while $\chi_{\text{top}} = 8$ for the surface $(F = 0)$ (with the obvious equality $\chi_{\text{top}}(X) = \chi_{\text{top}}(\tilde{X})$). In particular, since $[(G = 0)] = [\mathbb{P}^2] + 6\mathbb{L}$ and $[(F = 0)] = \mathbb{L}^2 + 4\mathbb{L} + 2[\mathbb{P}^1]$, both $X' := X \setminus \{\text{all singular fibers}\}$ and $\tilde{X}' := \tilde{X} \setminus \{\text{all singular fibers}\}$ satisfy $[X'] - [\tilde{X}'] = [X] - [\tilde{X}]$, with all the fibers on X' (resp. on $\tilde{X}'$) having the same class $(= [\mathbb{P}^2] + 6\mathbb{L})$ in $K_0(\text{Var})$. It is then tempting to propose that $[X'] = [\tilde{X}']$, but we don't have a rigorous argument for the proof, unfortunately.

(same for $\tilde{X}$) and [any smooth cubic surface] $= [\mathbb{P}^2]+6\mathbb{L}$, it suffices to consider $X_0, \tilde{X}_0$ in place of $X, \tilde{X}$, respectively.

Further, we have $\mathcal{X}_0 \simeq \tilde{\mathcal{X}}_0$, since one may perform a linear transformation of y_i (resp. of λ) which brings L_α to $L_{\tilde{\alpha}}$ (resp. $L_\beta + \lambda$ to $L_{\tilde{\beta}} + \lambda$). Then Lemma 2.4 yields $X_0 \times \mathbb{A}^{m+1} \simeq \tilde{X}_0 \times \mathbb{A}^{m+1}$ and so $[X_0] \cdot \mathbb{L}^{m+1} = [\tilde{X}_0] \cdot \mathbb{L}^{m+1}$. In the same way we obtain $[X_0] \cdot \mathbb{L}^m = [\tilde{X}_0] \cdot \mathbb{L}^m$ (using the second assertion of Lemma 2.4). The claim now follows by induction on the number of variables y_i in the linear forms $L_\alpha, L_{\tilde{\alpha}}$, etc. □

Corollary 2.8. $[X] \cdot \mathbb{L}^k \cdot (\mathbb{L}-1)^{m-k} = [\tilde{X}] \cdot \mathbb{L}^k \cdot (\mathbb{L}-1)^{m-k}$ *for all* $0 \le k < m$.

Proof. Again, it suffices to consider $X_0, \tilde{X}_0$ in place of $X, \tilde{X}$.

Set $L_i := \mathbb{A}^m \cap (y_i = 0), 1 \le i \le m$, and observe that $X_0 \times \mathbb{A}^k \times (\mathbb{A}^1 \setminus \{0\})^{m-k} \simeq X_0 \times \mathbb{A}^m \setminus \bigcup_{i=1}^{m-k} X_0 \times L_i$ (resp. $\tilde{X}_0 \times \mathbb{A}^k \times (\mathbb{A}^1 \setminus \{0\})^{m-k} \simeq \tilde{X}_0 \times \mathbb{A}^m \setminus \bigcup_{i=1}^{m-k} \tilde{X}_0 \times L_i$) for all $k < m$. Moreover, since $X_0 \times L_i \simeq \tilde{X}_0 \times L_i$ (see Proposition 2.7) and the schemes $X_0 \times L_i$ (resp. $\tilde{X}_0 \times L_i$) are the same for all i, the isomorphisms $X_0 \times L_i \simeq \tilde{X}_0 \times L_i$ glue together and give

$$X_0 \times \mathbb{A}^m \setminus \bigcup_{i=1}^{m-k} X_0 \times L_i \simeq \tilde{X}_0 \times \mathbb{A}^m \setminus \bigcup_{i=1}^{m-k} \tilde{X}_0 \times L_i. \tag{2.9}$$

Namely, each isomorphism $X_0 \times L_i \simeq \tilde{X}_0 \times L_i$ extends to that between $X_0 \times \mathbb{A}^m$ and $\tilde{X}_0 \times \mathbb{A}^m$, mapping y_i to y_i. Then, given any two $0 \le i, j < m$, we glue two copies of $X_0 \times \mathbb{A}^m$ (resp. of $\tilde{X}_0 \times \mathbb{A}^m$) over $X_0 \times \mathbb{A}^m \setminus L_i$ and $X_0 \times \mathbb{A}^m \setminus L_j$, respectively (same for $\tilde{X}_0$), where the gluing isomorphism just interchanges y_i and y_j. The so obtained variety contains $X_0 \times L_i \cup X_0 \times L_j$ whose complement coincides with $X_0 \times \mathbb{A}^m \setminus (X_0 \times L_i \cup X_0 \times L_j)$ (and similarly for $\tilde{X}_0$). We also have $X_0 \times \mathbb{A}^m \setminus (X_0 \times L_i \cup X_0 \times L_j) \simeq \tilde{X}_0 \times \mathbb{A}^m \setminus (\tilde{X}_0 \times L_i \cup \tilde{X}_0 \times L_j)$ by construction. Iterating this procedure yields (2.9).

The pertinent equalities now follow from the fact that

$$[X_0 \times \mathbb{A}^m \setminus \bigcup_{i=1}^{m-k} X_0 \times L_i] = [X_0] \cdot (\mathbb{L}^m - [\bigcup_{i=1}^{m-k} L_i]) = [X_0] \cdot \mathbb{L}^k \cdot (\mathbb{L}-1)^{m-k}$$

(same for $\tilde{X}_0$). □

Letting $k = 0$ in Corollary 2.8 we get the identity $[X] \cdot (\mathbb{L}-1)^m = [\tilde{X}] \cdot (\mathbb{L}-1)^m$. This together with Proposition 2.7 gives $[X] = [\tilde{X}]$ (we have expanded $(\mathbb{L}-1)^m = \mathbb{L}^m - m\mathbb{L}^{m-1} + \ldots + (-1)^m$ in (commutative) $K_0(\mathrm{Var})$ in the usual way).

2.10. Both X and $\tilde{X}$ satisfy the genericity assumption from [14, §1] concerning the pencils of degree 3 del Pezzo surfaces (cf. Lemma 2.3 and **2.5** above). Hence $X, \tilde{X}$ are not birational to each other (see [14, Corollary 2.1,(i)]). On the other hand, we have shown that $[X] = [\tilde{X}]$, and Theorem 1.2 follows.

Acknowledgments. I am grateful to A. Bondal, S. Galkin, M. Kapranov, V. Lunts, B. Poonen, A. Pukhlikov, and J. Sebag for their interest, helpful discussions and corrections. The work was supported by World Premier International Research Initiative (WPI), MEXT, Japan, and Grant-in-Aid for Scientific Research (26887009) from Japan Mathematical Society (Kakenhi).

References

[1] A. I. Bondal, M. Larsen and V. A. Lunts, Grothendieck ring of pretriangulated categories, Int. Math. Res. Not. **2004**, no. 29, 1461–1495.

[2] A. Chambert-Loir, J. Nicaise and J. Sebag, Motivic integration, to appear in Prog. in Math.

[3] A. Craw, An introduction to motivic integration, in *Strings and geometry*, 203–225, Clay Math. Proc., 3, Amer. Math. Soc., Providence, RI.

[4] J. Denef and F. Loeser, Germs of arcs on singular algebraic varieties and motivic integration, Invent. Math. **135** (1999), no. 1, 201–232.

[5] S. Galkin and E. Shinder, The Fano variety of lines and rationality problem for a cubic hypersurface, Preprint arXiv:1405.5154.

[6] M. Gromov, Endomorphisms of symbolic algebraic varieties, J. Eur. Math. Soc. (JEMS) **1** (1999), no. 2, 109–197.

[7] J. Kollár, Conics in the Grothendieck ring, Adv. Math. **198** (2005), no. 1, 27–35.

[8] M. Kontsevich. Motivic integration, Lecture at Orsay (1995).

[9] M. Larsen and V. A. Lunts, Motivic measures and stable birational geometry, Mosc. Math. J. **3** (2003), no. 1, 85–95, 259.

[10] M. Larsen and V. A. Lunts, Rationality of motivic zeta function and cut-and-paste problem, Preprint arXiv:1410.7099.

[11] E. Looijenga, Motivic measures, Astérisque No. 276 (2002), 267–297.

[12] M. Mustaţă, Jet schemes of locally complete intersection canonical singularities, Invent. Math. **145** (2001), no. 3, 397–424.

[13] B. Poonen, The Grothendieck ring of varieties is not a domain, Math. Res. Lett. **9** (2002), no. 4, 493–497.

[14] A. V. Pukhlikov, Izv. Ross. Akad. Nauk Ser. Mat. **62** (1998), no. 1, 123–164; translation in Izv. Math. **62** (1998), no. 1, 115–155.

Kavli IPMU (WPI), The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, 277-8583, Japan

E-mail address: ilya.karzhemanov@ipmu.jp